



	Known COVID-19 Positive	Unable to Screen	Screened Positive (Symptomatic per SHA Screening tool)*	Screened Positive <b>ASYMPTOMATIC</b> (Travel/Contact/Mass Gathering History or from Area of <b>Elevated Activity</b> )	ASYMPTOMATIC and Screened Negative
<b>Decision</b>	<ul style="list-style-type: none"> <li>• Droplet/Contact Plus precautions</li> </ul>	<ul style="list-style-type: none"> <li>• Droplet/Contact Plus precautions until able to screen</li> </ul>	<ul style="list-style-type: none"> <li>• Droplet/Contact Plus precautions</li> </ul>	<ul style="list-style-type: none"> <li>• Droplet/Contact Plus precautions</li> <li>• monitor for COVID-19 symptoms</li> </ul>	<ul style="list-style-type: none"> <li>• Routine precautions</li> </ul>
<b>Testing if no previous swab done</b>	<ul style="list-style-type: none"> <li>• Not required</li> </ul>	<ul style="list-style-type: none"> <li>• Swab for COVID-19 based on provincial testing guidelines</li> </ul>	<ul style="list-style-type: none"> <li>• Swab for COVID-19 based on provincial testing guidelines</li> </ul>	<ul style="list-style-type: none"> <li>• Offer testing per provincial admission swab guidelines</li> <li>• Not required unless symptoms develop</li> </ul>	<ul style="list-style-type: none"> <li>• Not mandatory</li> <li>• Offer testing as per provincial admission swab guidelines</li> </ul>
<b>Testing if admission swab previously collected and negative</b>	<ul style="list-style-type: none"> <li>• Not required</li> </ul>	<ul style="list-style-type: none"> <li>• Repeat swab not indicated unless clinical suspicion of COVID-19</li> </ul>	<ul style="list-style-type: none"> <li>• If symptoms worsen, repeat swab 48 hours after symptom onset OR 24 hours after initial test, whichever is greater</li> </ul>	<ul style="list-style-type: none"> <li>• If symptoms develop, swab 48 hours after symptom onset OR 24 hours after initial test, whichever is greater</li> </ul>	<ul style="list-style-type: none"> <li>• Not required</li> </ul>
<b>COVID-19 positive facility maternity care?</b>	<b>Yes</b>	<ul style="list-style-type: none"> <li>• Notify anesthesia and pediatrics/newborn care team</li> <li>• Consult obstetrics if not under Obstetrical service</li> </ul>		<ul style="list-style-type: none"> <li>• Notify anesthesia, obstetrics and pediatrics/ newborn care team</li> </ul>	<ul style="list-style-type: none"> <li>• Routine care</li> </ul>
	<b>No</b>	<ul style="list-style-type: none"> <li>• Transfer to facility planned for COVID-19 positive maternity care if possible</li> <li>• If transfer not feasible, implement <a href="#">Intrapartum Care Standard for COVID-19 (CV-19 WS0021)</a></li> <li>• Notify available supporting care teams</li> </ul>		<ul style="list-style-type: none"> <li>• <b>Deliver in planned delivery site</b></li> </ul>	<ul style="list-style-type: none"> <li>• Deliver in planned delivery site</li> </ul>



	Known COVID-19 Positive	Unable to Screen	Screened Positive (Symptomatic per SHA Screening tool)*	Screened Positive <b>ASYMPTOMATIC</b> (Travel/Contact/Mass Gathering History or from Area of <b>Elevated Activity</b> )	ASYMPTOMATIC and Screened Negative
<b>C-section</b>	<ul style="list-style-type: none"> <li>Follow <a href="#">CV-19 G0005 Protocol for Operative Management of Surgical Patients (All Ages) during COVID-19</a>.</li> <li>Regional anesthetic – utilize and optimize whenever possible                             <ul style="list-style-type: none"> <li>Surgeon and anesthesiologist in <b>full N95 PPE</b> in case conversion to GA required</li> <li>Remainder of team in surgical masks / routine precautions</li> <li>Leave theatre to doff and don <b>full N95 PPE</b> if conversion to GA required</li> <li>Theatre specific settle time is not possible after intubation</li> </ul> </li> <li>Planned General anesthetic                             <ul style="list-style-type: none"> <li>Whole team in <b>full N95 PPE</b> (Fit-tested N95 respirator; Face shield or goggles)</li> <li>Theatre specific settle time is not possible after intubation</li> </ul> </li> </ul>				<ul style="list-style-type: none"> <li>Utilize regional anesthetic whenever possible</li> <li>Identical process for conversion of regional to general anesthesia</li> <li>General anesthesia routine precautions for intubation/extubation</li> </ul>
<b>Modification of precautions***</b>	<ul style="list-style-type: none"> <li>May discontinue precautions 14 days after positive swab or symptom onset, or 48 hours after resolution of symptoms (whichever is later)</li> <li>No retesting for COVID-19 required</li> </ul>	<ul style="list-style-type: none"> <li>Precautions may be lifted once:                             <ul style="list-style-type: none"> <li>screening can be done and is <b>NEGATIVE</b> <u>AND</u></li> <li>Swab result is negative</li> </ul> </li> </ul>	<p><b>If FEVER ALONE</b> is the only reason for positive screen, precautions not required IF:</p> <ul style="list-style-type: none"> <li>Swab result is negative <u>AND</u></li> <li>Alternate source of fever identified</li> </ul> <p><b>FEVER and/or other ILI symptoms:</b></p> <ul style="list-style-type: none"> <li>Follow algorithm <a href="#">CV-19 A0002 for COVID-19 Negative Inpatients and Residents</a></li> </ul>	<ul style="list-style-type: none"> <li>Continue Contact/droplet plus precautions for 14 days after last exposure</li> </ul>	<ul style="list-style-type: none"> <li>Not required</li> </ul>

\* Refers to [SHA 0002 Screening Tool](#). COVID-19 should be considered high in the differential for unexplained intrapartum fever and/or cough. "Aches and pains" refer to flu-like symptoms and not labour pains. This tool is not intended to replace clinical judgement in individual patient management and alternate diagnosis must be considered before the patient's final risk of COVID-19 is determined.

\*\* Retesting is done only for epidemiological reasons.

\*\*\*Removal/step down of precautions should only be made in consultation with the most responsible physician (MRP).