



**INFORMED CONSENT**

- **Currently, there is no information on administering a COVID-19 vaccine and another vaccine at the same time.**
- **Administration of the COVID-19 vaccine close together with another vaccine may potentially result in lessened immune response to the vaccine or mistakenly attribute an adverse event to a particular vaccine.**
- **Based on your immunization history, it is preferable to wait at least 14 days to receive the COVID-19 vaccine.**
- **Do you have any additional questions or concerns about getting immunized with the COVID-19 vaccine?**

SCENARIO	CONSENT	OPTIONS	RECOMMENDATION
Client presents at a COVID-19 vaccine clinic and has received any vaccine less than two weeks ago. (between 13 days to 1 day ago)	Provide Informed Consent	Client can rebook or represent two weeks after non-COVID vaccine.	Vaccinate with COVID vaccine as client is here and we have the opportunity to vaccinate
		Client can request vaccination.	
Client is pregnant and due for her Tdap vaccine. She presents for her COVID-19 vaccine.	Provide Informed Consent		Vaccinate with a COVID-19 vaccine and encourage her to contact Public health/HCP for a Tdap
A pregnant client calls to state she needs her Tdap vaccine, but received her COVID-19 vaccine 5 days ago.	Provide Informed Consent	Client is early in her eligibility (27 weeks) and can wait four weeks post COVID vaccine + have sufficient time before birth for optimal transfer of maternal antibodies (4 weeks).	Book closer to the four week interval
		Client has a history of early deliveries, has a high risk pregnancy for potential of early delivery, or is further along in her pregnancy.	Book Tdap
Client calls public health to book a routine vaccine and has received COVID-19 less than 4 weeks ago.	Provide Informed Consent	Is there an outbreak	Book vaccine
		Client due for vaccine.	Book based on preferred interval of 4 weeks from COVID-19 vaccine.
Client calls public health to book a vaccine that falls under a special population. <ul style="list-style-type: none"> <li>• Td for injury</li> <li>• Cocooning</li> <li>• Blood and body fluid exposure</li> <li>• Rabies</li> <li>• Immunocompromised</li> <li>• Post exposure vaccine</li> </ul>	Provide Informed Consent	Client has a current risk for transmitting or obtaining a Non-COVID disease.	Book in for vaccine irrespective of 4 week interval
		Client's risk is in the near future (i.e. cocooning and baby will not be born for 6 weeks)	Potential risk in the future. Book appointment honoring the preferred interval of four weeks.